注意:考試開始鈴響前,不得翻閱試題,並不得書寫、畫記、作答。

國立清華大學 114 學年度碩士班考試入學試題

系所班組別:生命科學暨醫學院

丁組(醫學生物科技學程)

科目代碼:0701

考試科目:生物化學

一作答注意事項-

- 1. 請核對答案卷(卡)上之准考證號、科目名稱是否正確。
- 考試開始後,請於作答前先翻閱整份試題,是否有污損或試題印刷不清,得舉手請監試人員處理,但不得要求解釋題意。
- 3. 考生限在答案卷上標記 由此開始作答」區內作答,且不可書寫姓 名、准考證號或與作答無關之其他文字或符號。
- 4. 答案卷用盡不得要求加頁。
- 5. 答案卷可用任何書寫工具作答,惟為方便閱卷辨識,請儘量使用藍色或黑色書寫;答案卡限用 2B 鉛筆畫記;如畫記不清(含未依範例畫記)致光學閱讀機無法辨識答案者,其後果一律由考生自行負責。
- 6. 其他應考規則、違規處理及扣分方式,請自行詳閱准考證明上「國立 清華大學試場規則及違規處理辦法」,無法因本試題封面作答注意事項 中未列明而稱未知悉。

系所班組別:生命科學暨醫學院甲組、乙組、丁組

·考試科目(代碼):生物化學(0401、0501、0701)

共_7_頁,第_1_頁 *請在【答案卡】作答

Part 1 選擇題, 20 題單選題 (每題 1 分, 共 20 分), 答錯不倒扣

- 1. Which of the following reaction would favor cellular gluconeogenesis in the fasted state?
 - (A) Acetyl-CoA activates pyruvate carboxylase.
 - (B) Fructose 2,6-bisphosphateactivates pyruvate kinase.
 - (C) AMP activates phosphfructokinase 1.
 - (D) NAD⁺ activates glyceraldehyde 3-phosphate dehydrogenase.
 - (E) None of the above.
- 2. Which of the following urine component can be used as a biomarker to monitor DNA turnover?
 - (A) beta-alanine.
 - (B) ammonium ion
 - (C) beta-aminoisobutyrate.
 - (D) cytidine.
 - (E) AMP
- 3. Which of the following enzyme is not located in mitochondria?
 - (A) carbamoyl-phosphate synthase 1
 - (B) voltage-dependent anion channel
 - (C) carnitine transporter
 - (D) DNA polymerase gamma
 - (E) carbamoyl-phosphate synthase 2
- 4. Which inhibitor can be used to inhibit complex III of electron transport chain in cells?
 - (A) cyanide
 - (B) oligomycin
 - (C) rotenone
 - (D) malonate
 - (E) none of the above

系所班組別:生命科學暨醫學院甲組、乙組、丁組

考試科目(代碼):生物化學(0401、0501、0701)

共_7_頁,第_2_頁 *請在【答案卡】作答

- 5. Which of the following best describes the relationship between glucose and fructose?
 - (A) Epimers
 - (B) Structural isomers
 - (C) Enantiomers
 - (D) Diastereomers
 - (E) Cis-trans isomers
- 6. What important reducing agent is an important product of the pentose phosphate pathway?
 - (A) NADH
 - (B) NADPH
 - (C) NADP+
 - (D) NAD+
 - (E) DTT
- 7. Which glycolytic reaction cannot be directly reversed in gluconeogenesis?
 - (A) glucose converted into glucose-6-phosphate
 - (B) glucose-6-phosphate converted into fructose-6-phosphate
 - (C) 3-phosphoglycerate converted into 1,3-bisphosphoglycerate
 - (D) 1,3-bisphosphoglycerate converted into glyceraldehyde-3-phosphate
 - (E) phosphoenolpyruvate converted into 2-phosphoglycerate
- 8. Which of the following is catalyzed by an oxidoreductase in Krebs cycle?
 - (A) oxaloacetate > citrate
 - (B) isocitric acid > α-ketoglutaric acid
 - (C) fumarate > malate
 - (D) citric acid > isocitric acid
 - (E) citric acid > cis-aconitate
- 9. Why is single-stranded RNA (ssRNA) chemically less stable than single-stranded DNA (ssDNA)?
 - (A) ssRNA has a more rigid structure than ssDNA.
 - (B) because Uridine is less stable than thymidine

系所班組別:生命科學暨醫學院甲組、乙組、丁組

考試科目(代碼):生物化學(0401、0501、0701)

共_7_頁,第_3_頁 *請在【答案卡】作答

(C)	ssDNA	is more	flexible	than	ssRNA.
-----	-------	---------	----------	------	--------

- (D) ssRNA has stronger base stacking interactions than ssDNA.
- (E) ssRNA contains an additional hydroxyl group that makes it more reactive.
- 10. Which amino acid is the most flexible in protein structures?
 - (A) Alanine
 - (B) Leucine
 - (C) Proline
 - (D) Serine
 - (E) Glycine
- 11. Flippases are enzymes that flip:
 - (A) fatty acids from one site on glycerol to another.
 - (B) glucose from its alpha- to beta-configuration.
 - (C) amino acids between different positions within a protein.
 - (D) cholesterol between different organelles.
 - (E) phospholipids to the opposite side of a membrane.
- 12. How many chiral carbons are present in an aldopentose?
 - (A) 1 (B) 2
- (C) 3
- (D) 4
- (E)5
- 13. Aminotransferases are enzymes that have all of the following characteristics except:
 - (A) Can make a α-keto acid from an α-amino acid
 - (B) use pyridoxal phosphate as a carrier of amino acid
 - (C) catalyze highly reversible reactions
 - (D) Use ATP
 - (E) Serum levels are used as a clinical measure of tissue damage
- 14. Pyruvate can be converted to glucose via gluconeogenesis or oxidized to acetyl-CoA.
 - (A) In the conversion of pyruvate to glucose, the first enzyme pyruvate carboxylase is allosterically inhibited by acetyl-CoA.

系所班組別:生命科學暨醫學院甲組、乙組、丁組

考試科目(代碼):生物化學(0401、0501、0701)

共_7_頁,第_4_頁 *請在【答案卡】作答

- (B) In the conversion of pyruvate to glucose, the first enzyme pyruvate dehydrogenase is allosterically inhibited by acetyl-CoA.
- (C) Pyruvate is oxidized to acetyl-CoA by pyruvate dehydrogenase complex, which is allosterically inhibited by acetyl-CoA.
- (D) Pyruvate is oxidized to acetyl-CoA by pyruvate dehydrogenase complex, which is allosterically activated by acetyl-CoA.
- (E) None of the above.
- 15. In the citric acid cycle, all of the following enzymes require NAD+ as a coenzyme except the reaction catalyzed by:
 - (A) isocitrate dehydrogenase.
 - (B) malate dehydrogenase.
 - (C) a-Ketoglutarate dehydrogenase.
 - (D) succinate dehydrogenase.
 - (E) none of the above.
- 16. HMG-CoA reductase is the key regulatory site in cholesterol synthesis. Which of the following statement about the regulatory mechanisms of HMG-CoA reductase is TRUE?
 - (A) HMG-CoA reductase phosphatase can activate HMG-CoA reductase,
 - (B) A high AMP:ATP ratio leads to activation of HMG-CoA reductase,
 - (C) Phosphorylation by cAMP-dependent kinase activates HMG-CoA reductase,
 - (D) High cholesterol level increases the half-life of HMG-CoA reductase,
 - (E) High cholesterol level increases HMG-CoA reductase gene expression.
- 17. What is the role of transition-state stabilization in enzyme-catalyzed reactions?
 - (A) To enhance the thermal stability of the enzyme
 - (B) To increase the binding affinity of the substrate
 - (C) To lower the activation energy, thereby increasing the reaction rate
 - (D) To increase the stability of the product, thereby reducing side reactions
 - (E) To provide reaction versatility

系所班組別:生命科學暨醫學院甲組、乙組、丁組

考試科目(代碼):生物化學(0401、0501、0701)

共_7_頁,第_5_頁 *請在【答案卡】作答

- 18. What is the correct order of these events during muscle contraction?
 - a. Calcium ions are released from the sarcoplasmic reticulum
 - b. Hydrolysis of ATP and energy release
 - c. Detachment of myosin from actin
 - d. Power stroke
 - e. Opening of switch 1 and switch 2 on myosin head
 - (A) a e b d c (B) e b c a d (C) a b c d e (D) a d e b c (E) e d c b a
- 19. Enzyme A can act on substrate S and substrate T and follows simple Michaelis-Menten kinetics. The Michaelis constant K_m of enzyme A for substrate S is $K_m^S = 2$ mM, and the rate constant k^S is $5X10^3$ sec⁻¹. For substrate T, $K_m^T = 5$ mM, and k^T is $2.5X10^4$ sec⁻¹. Which substrate exhibits greater catalytic efficiency?
 - (A) S
 - (B) T
 - (C) Enzyme A uses S and T with the same catalytic efficiency.
 - (D) cannot be determined due to insufficient information.
 - (E) The result would change if the enzyme's conditions or substrate concentrations varied.
- 20. Hemoglobin exhibits sigmoidal binding kinetics for oxygen. Which of the following terms describes hemoglobin's ability to enhance oxygen binding after the first oxygen molecule binds?
 - (A) Allosteric inhibition
 - (B) Competitive inhibition
 - (C) Allosteric activation
 - (D) Cooperativity
 - (E) Catalysis

系所班組別:生命科學暨醫學院甲組、乙組、丁組

考試科目(代碼):生物化學(0401、0501、0701)

共_7_頁,第_6_頁 *請在【答案卷】作答

Part 2 簡答題 10 題 (每題 8 分, 共 80 分)

- 1. Why do polysaccharides have greater structural diversity compared to polypeptides? (8 points)
- Which two amino acids are completely positively charged at physiological pH? (8 points)
- 3. Please describe or illustrate the structure (2 points) of ATP synthase and how this machinery links to the chemiosmotic hypothesis (6 points).
- 4. Please describe how starch is synthesized in plants (4 points) and how this process is regulated by photosynthesis (4 points).
- 5. How is the organic sulfide assimilated from sulfate? (8 points)
- 6. Please compare the carbamoyl-phosphate I and II according to their substrate; intracellular locations; involved pathways; regulators. (8 points)
- 7. How are eukaryotic transcripts processed and delivered to the ribosomes for translation? (8 points)
- 8. What is the structure of ribosomes, and how are they assembled? (8 points)
- Please design a set of oligonucleotide primers (16-mers) of the following short DNA sequence for PCR amplification (5' -> 3'). (8 points)
 TGTGGGAGCTGTGATGATGTTTGTAGGCTTCCTGGGGTGCTATGGGGCCATC CAGGAGGTGGCTGCAGGCATCTGGGGCTTCGTAAACAAAGACCAGATCGC CAAGGA

系所班組別:生命科學暨醫學院甲組、乙組、丁組

考試科目(代碼):生物化學(0401、0501、0701)

共_7_頁,第_7_頁 *請在【答案卷】作答

- 10. A 12-kb DNA fragment was digested with restriction enzymes HindIII and PstI under the following conditions:
 - (1) Digestion with HindIII yielded fragments of 3 kb, 4 kb, and 5 kb.
 - (2) Digestion with PstI yielded fragments of 2 kb, 5 kb, and 5 kb.
 - (3) Concomitant digestion with both HindIII and PstI yielded fragments of 1 kb, 2 kb, 2 kb, 3 kb, and 4 kb.

Using this data, please construct a possible restriction map of the original 12-kb fragment. (8 points)