

國立清華大學 100 學年度碩士班入學考試試題

系所班組別：核子工程與科學研究所甲組(工程組)

考試科目 (代碼)：工程數學(3001)


共 3 頁，第 1 頁 *請在【答案卷、卡】作答

1. Find the transient current if

$$R = 6 \Omega, L = 1 H, C = 0.04 F, E = 600 (\cos t + 4 \sin t) V;$$

(L, R, C, E, are measured in henrys, ohms, farads, volts, respectively.)

Initial current and charge are assumed to be zero. (10%)



2. Consider a general nonhomogeneous linear ODE

$$y^{(n)} + p_{n-1}(x)y^{(n-1)} + p_{n-2}(x)y^{(n-2)} + \dots + p_1(x)y' + p_0(x)y = r(x)$$

The particular solution $y_p(x)$ can be solved by the method of variation of parameters. That is,

$$y_p(x) = \sum_{k=1}^n y_k(x) \int \frac{W_k(x)}{W(x)} r(x) dx$$

(Where the y_k 's are n linearly independent homogeneous solutions. W is the Wronskian of y_1, \dots, y_n , and W_k is identical to W, but with the kth column replaced by a column of zeros-except for the bottom element, which is 1.)

Try to solve the following equation using the method of variation of parameters.

$$y''' + \frac{3}{4}x^{-2}y' - \frac{3}{4}x^{-3}y = 9x^{5/2} \quad (10\%)$$

國立清華大學 100 學年度碩士班入學考試試題

系所班組別：核子工程與科學研究所甲組(工程組)

考試科目 (代碼)：工程數學(3001)

共 3 頁，第 2 頁 *請在【答案卷、卡】作答

3. Solve the following equation by Laplace Transform method.

$$y' + y = f(t), \quad y(0) = 3,$$
$$\text{where } f(t) = \begin{cases} 0 & 0 \leq t < \pi \\ 2 \cos t & t \geq \pi \end{cases} \quad (10\%)$$

4. Find the basis of solutions $y(x)$ of the following differential equation. Show the details of your work.

$$xy'' + (2x + 1)y' + (x + 1)y = 0. \quad (10\%)$$

5. Find a unit vector normal to surface S given by $\cos(xy) = e^z - 1$ at the point $(1, \pi, 0)$. (10%)

6. Let $\mathbf{F} = (x-y)\mathbf{i} + (y-z)\mathbf{j} + (z-x)\mathbf{k}$. Evaluate the surface integral of \mathbf{F} over the unit sphere defined by $x^2 + y^2 + z^2 = 1$. (10%)

7. Define the Fourier transform of $f(x)$ to be $\hat{f}(w) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x)e^{-ixw} dx$.

(a) [5%] Calculate the Fourier transform of $f(x) = \begin{cases} a - |x| & , |x| < a \\ 0 & , \text{otherwise} \end{cases}$

(b) [5%] Consider the one-dimensional diffusion equation:

$$\frac{\partial}{\partial t} u(x, t) = D \frac{\partial^2}{\partial x^2} u(x, t) \quad \text{for } -\infty < x < \infty$$

with the initial condition $u(x, 0) = f(x)$. Use the Fourier transform to show that the

solution of the diffusion equation takes the form $u(x, t) = \int_{-\infty}^{\infty} K(x - \xi, t) f(\xi) d\xi$.

Find $K(x - \xi, t)$, which is called the kernel.

[Hint: Gaussian integral $\int_{-\infty}^{\infty} \exp\left(\frac{-x^2}{2\sigma^2}\right) dx = \sqrt{2\pi\sigma^2}$]

國立清華大學 100 學年度碩士班入學考試試題

系所班組別：核子工程與科學研究所甲組(工程組)

考試科目 (代碼)：工程數學(3001)

共 3 頁，第 3 頁 *請在【答案卷、卡】作答

8. (a) [5%] Find and classify all local maxima, local minima and saddles for

$$f(x, y, z) = \exp(2x^2 + xz - 5z^2).$$

(b) [5%] Consider a forced vibration system which is described by the equations

$$\begin{aligned} \frac{d^2 x_1}{dt^2} + 2x_1 - x_2 &= A \sin(\omega t) \\ \frac{d^2 x_2}{dt^2} - x_1 + 2x_2 &= B \sin(\omega t) \end{aligned}, \text{ where } A, B, \text{ and } \omega \text{ are constant.}$$

To seek a particular solution, we assume $x_1(t) = q_1 \sin(\omega t)$ and $x_2(t) = q_2 \sin(\omega t)$. Find q_1 and q_2 .

9. Along the circumference of the circle $r = b$ a solution $T(r, \theta)$ of Laplace's equation is required to take on the value T_0 when $0 < \theta < \pi$ and the value $-T_0$ when $\pi < \theta < 2\pi$. Determine an expression for T valid when $r > b$. (Show the details of your work.)

$$[\nabla^2 T = \frac{1}{r} \frac{\partial}{\partial r} (r \frac{\partial T}{\partial r}) + \frac{1}{r^2} \frac{\partial^2 T}{\partial \theta^2}] \quad (12\%)$$

10. (complex analysis) Prove the following identity

$$\sin^{-1} z + \cos^{-1} z = \frac{1}{2}(4n+1)\pi, \quad n = 0, \pm 1, \pm 2, \dots \quad (8\%)$$